ANALYSIS OF FACTORS RELATED TO IMPLEMENTATION OF COMPLETE BASIC IMMUNIZATION IN WORK AREA OF PUSKESMAS PENGAMBIRAN PADANG CITY

Ade Nurhasanah Amir*

Department of Midwife Science, Syedza Saintika Institute of Health Science

* Corresponding author : Adhek nurhasanahamir@yahoo.com

ABSTRACT

The role of a mother in the immunization program is very important, so the understanding of immunization is necessary, the behavior of a person or society about health is determined by knowledge, attitude, belief, tradition, work etc. from the person or society concerned. In addition, the availability of health care facilities, attitudes, behavior of health workers on the provision of health information will also support and strengthen the formation of health behaviors, This study aims to determine the factors that affect the provision of complete basic immunization, This research uses a combined research method between quantitative and qualitative. Research sample for quantitative research is 107 mother mother, while informant for qualitative research is Head of Puskesmas Pengambiran Padang City, Responsible, Immunization Officer of Puskesmas and Kader which is in Working Area of Pengambiran Padang City, The result of research is obtained the factor that play the most role in the implementation Complete basic immunization is family support (p = 0.001). The results of the analysis revealed that the coverage of the implementation of complete basic immunization has not been successful because the lack of active role of cadres and health officers in the midst of the community in informing immunization in depth so that there is still a lack of understanding from mothers and families which no family support in the implementation of complete basic immunization, has not reached 95% target because the system implementation of complete basic immunization services that have not been optimal. It needs various efforts to improve the achievement of complete basic immunization coverage especially through optimizing the role and function of health and cadre officers and optimizing partnerships with various parties

Keyword: Basic Immunization, Toddler

INTRODUCTION

Health development aims to improve public health status in order to improve the quality of life, intelligence and community welfare (1). This is then outlined in the Millennium Development Goals (MDGs) which is a global and national commitment to improve the welfare of the people, one of the goals of the MDGs is to reduce child mortality with a target of reducing child mortality under five years by two-thirds in the period from 1990 to 2015 means that it decreases from 97 per 1000 live births to 32 per 1000 live births (1). The fact that

the level of health is still low especially in the poor reflected from the high AKI and IMR, this is influenced by the lack of clean and healthy life behavior has not been entrenched in people who have knowledge and education is still low².

Efforts to reduce infant and under-five mortality include improving healthy behaviors, as well as concerns about child survival and early development, and increasing immunization coverage (3). Accelerating the decrease of under-five mortality rate by government is one of them is immunization program which is



promotive and preventive activity that have important role in decreasing infant and child mortality ⁴.

Over the past few years, concerns about the return of some infectious diseases and the rise of new infectious diseases are increasing (5). (30%), the incidence of malnutrition, not 100% of UCI National Village, the incidence of measles, varicella and polio cases. With the focus of intervention, among others, Quality Improvement and of health Ouantity personnel expected to decrease the number Neonatal death ²

The success of infants obtaining five basic immunization types (HB0, BCG, DPT-HB, Polio, and Measles) was measured through a complete basic immunization indicator, compared to the 2008-2011 period, the of coverage complete basic immunization during 2012-2015 Indonesia decreased (6). The full coverage of basic immunization based on routine data in 2010-2013 reaches the target of the Ministry of Health's Strategic Plan (Renstra), but by 2014 and 2015 immunization coverage does not reach the target of the Renstra expected⁶. There is still a perception in the community and even in the household that infant immunization and weighing are not important to be a challenge to the success of the health program ². Research in Bandung shows that there is relationship of mother characteristic of under five (63,4%) and mother perception of toddler immunization (64,8%) to complete basic immunization implementation ⁷.

The number of infant deaths in West Sumatra Province as many as 681 people spread in 19 districts / cities with the highest death contributor from the city of Padang as many as 108 people. The number of deaths of children under five in West Sumatera are 856 people

spread in 19 districts / cities with the highest death contributor from Padang City as many as 125 people (8). Research in the field of 2017 shows high infant mortality caused by several factors: maternal factors, infant factors, and health services. Maternal factors include social, low education, knowledge of health services and low economic levels, age less than 20 years or over 35 years. Toddlers, especially infants are a group of highly vulnerable populations with infections and disease attacks due to the development of organs and immune systems that have not been maximized. This condition causes many babies to die from illnesses that are not handled properly.1-3 Infant mortality refers to the death of a child under one year of age (9).

Baby target 0 - 11 months in 2014 is 102,040 babies. Achievement of Immunization program of west sumatera province in 2014 there are still some that have not been reached (8). Program Outreach is seen from first contact immunization coverage, Hb0: 86.62%, DPT-HB3: BCG: 92.63%, 86%. Meanwhile, the first contact target in 2014 is 95%. When compared to the achievement with the first contact target, no one has achieved the target, it is because there are still some people in community that illegal myth immunization, So for West Sumatera province in 2014 coverage complete basic immunization is still under the target of only 84.51% (8).

Based on the preliminary survey data conducted at the Padang City Health Office, the work area of puskesmas pengambiran is one of the lowest immunization coverage of 22 puskesmas in padang city, of 792 immunized infants got 66.6 percent (DKK, 2015). Based on these data it can be seen that Puskesmas Pengambiran is the village with the lowest immunization



coverage in 2015 with coverage of each immunization type as follows DPTHB1 (77,4%), DPT-HB2 (78,6%), DPT-HB3 (78, 0%), polio4 (73.3%), and measles (71.2%).Based these on immunization coverage generally has not fulfilled the target of $\geq 80\%$ ie DPT-HB, polio and measles (9). So from that research interested to examine factor factor what related completeness of basic immunization at baby in puskesmas pengambiran 2017 Based on the above description, the researcher is interested in taking the title of "Analysis of Factors related to Complete Basic Immunization of Infants in the Working Area of Puskesmas Pengambiran 2017. The purpose of this research is to analyze Factors related to the implementation of complete basic immunization in the region work of puskesmas in Padang City.

MATERIAL AND METHODS

The type of this research is combination research that is research method that combine quantitative research method and qualitative research method (9). Quantitative research uses sectional design. Data cross independent variables and dependent variables are collected and assessed at a Then qualitative research is conducted to dig deep information about the effort complete basic immunization management the for implementation of complete immunization in infant year 2017. This research was conducted in Working Puskesmas Pengambiran Area of Padang City. with research time in July-November 2017. The population in this study is all mothers who have under-fives under 2 years in the working Pengambiran Puskesmas which is about 292 people. Sampling was taken with multistage random sampling technique. Multistage random sampling technique is a sample selection technique that is done in stages and usually based on the division of work a government. Inclusion area of criteria Infant mothers who have KMS immunization records in infants, mothers who have babies 12-24 months, Mothers willing to be respondents, the number of research samples as many as 107 people. The data were collected by interview using questioner and observation / study of KIA book univaiat, bivariate document, and multivariate analyzed data.

RESULT

Puskesmas in Padang as many as 22 puskesmas. Puskesmas Pengambiran is one of the community health centers with low coverage of basic immunization with only 60% (8%), Batuang Village taba. The working area of Pengambiran Public Health Center is 27.21 km2 with the northern boundary with Tanjung saba, the south is bungus bay of sack, east of Lubuk refinery in the east, South Padang subdistrict (8)

Facilities and infrastructure owned by Puskesmas Pengambiran consisted of 1 primary health center, 1 sub health center, 4 units of poskeskel, 1 unit of mobile health center, 46 posyandu balita, 10 posyandu elderly, 187 cadres, 6 private doctors practice, 17 practice of private midwife (8). While there are 17 Posyandu bailita, 1 posyandu elderly, 7 practice midwives, private maternity hospital, in Pampangan sub-district there are 9 posyandu balita, 2 posyandu elderly, 4 independent midwives, 1 private practice physicians, Batuang Taba urban village have 9 posyandu balita, 3 posyandu elderly, 3 midwives of independent practice, and 1 maternity house, in kampuang jua village there are posyandu balita as many as 4 pieces, 2 posyandu elderly, 3



midwives of independent practice, and 1 maternity Hospital. and lastly in Kelurahan Gates there are posyandu

toddlers as many as 7 pieces, posyandu elderly 2 fruit, and maternity house 2 pieces (8).

a) Univariat Analysis

Dependen Variabels

Table 1
Frequency Distribution of Respondents Implementation of Complete Basic
Immunization in Working Area of Puskesmas Pengambiran Padang City Year 2017

Basic Immunization	on	F	%
Incomplete	53		49,5
Complete	54		50,5
Total	107		100

Based on table 5.1 it is seen that 53 respondents (49.5%) of respondents did not carry out

complete basic immunization in Puskesmas Pengambiran Padang working area.

b) Bivariat Analisis

Independent Variables

Table 2
Frequency Distribution of Respondents in the Working Area of Puskesmas
Pengambiran Padang City 2017

1 011 guilloit un 1 uuming 010, 2017				
	f	%		
attitude		·		
- Negatif	42	39,3		
- Positif	65	60,7		
Family Support				
- Not	56	52,3		
- There is	51	47,7		
Health Services				
- Not good	54	48,2		
- Well	53	47,3		

From the table it can be seen that more than half of the respondents

have a positive attitude, more than half do not get family support. and less than half get poor servic

c) Multivariat Analisis

Table 3
Most Dominant Related Factor Analysis Table with Complete Basic Immunization
Implementation

Variabel	В	SE	Wald	Sig	Ekp β
Step 1					
Health services	2,103	,913	5,301	,021	8,174
attitude	1,903	,605	9,907	,002	6,709

Oral Presentation



THE 1ST SYEDZA SAINTIKA INTERNATIONAL CONFERENCE ON NURSING, MIDWIFERY, MEDICAL LABORATORY TECHNOLOGY, PUBLIC HEALTH, AND HEALTH INFORMATION MANAGEMENT (SeSICNIMPH)

Family support 7,944 2,088 12,033 ,001 8,187

To see which variable has the greatest influence on the independent variable, look at exp (B). The greater the value of exp (B), the greater the effect on the dependent variable. The value of the betha exponent (exp B) is the greatest in the family support variable, which is 8.187, meaning that if family support can be implemented properly, the implementation of complete basic immunization is 8 times better.

Qualitative Research Results

d) Characteristics of Informants

Oualitative data collection conducted using in-depth interviews and focus group discussions (FGDs). In-depth interviews were conducted to the heads of puskesmas, immunization managers, immunization officers, cadres. FGD Conducted on 10 Mothers who have children under five in the working area Puskesmas Pengambiran Padang. was conducted respondents who had previously filled out the questionnaire in this study. Informants in this study amounted to 14 people.

Table 4
Characteristics of Informants in Implementation of Complete Basic Implementation in
Working Area of Puskesmas Pengambiran Kota Padang

Informan	Initial	Age	Backgroud	Education	Method
1	DN	40	Health Of Public center	S1	
2	S	30	Public center	S1	Depth
3	ES	33	Public center	S1	interview
4	EA	40	Kader	SMA	
5	RK	35	Mother	SMA	
6	E	21	Mother	SMA	
7	RS	22	Mother	SMA	
8	J	28	Mother	SMA	
9	S	34	Mother	SMA	
10	SI	29	Mother	SMA	FGD
11	ANP	31	Mother	SMA	
12	ANY	35	Mother	SMA	
13	DK	27	Mother	SMA	
14	AJ	27	Mother	SMA	

DISCUSSION

1. Attitude

Based on the result of the research, most of the respondents of the implementation of basic immunization have negatife attitude.

The results are in line with the results of research Paridawati (2013). In the study showed that attitudes have a role to the basic immunization.

The study findings are not in line with the research of Febriana (2009), the research shows that the



attitude does not affect the completeness of basic immunization. Approximately 81.58% of parents belong to a good attitude, but the good attitude is not followed by the high number of immunization completeness which only reached 46.78%.

Based on FGD results and in-depth interviews with respondents who have children with an incomplete immunization status who are mostly sociocultural or that assume that beliefs immunization is unnecessary. Respondents with socioculture or beliefs tend to have negate attitudes towards immunization, so that many of the immunization status of their children is incomplete and does not Besides immunize. misunderstanding of the community about the side effects immunization also greatly affect the attitude of the community against immunization (12)

Attitude is an internal reaction of a person who influenced by various factors of personal experience, culture, other important people, religion emotional factors in the individual who has an important role in the formation of attitudes. The process of attitudes caused by stimuli such as public knowledge. Such stimuli stimulate society to respon in the form of positive attitudes and negative attitudes that will eventually be realized in the form of concrete actions (13)

Attitudes consist of 4 levels of receiving, individuals wanting and paying attention to stimuli given, responding (responding), individual attitudes can provide answers when asked, doing and

completing tasks assigned, valuing; the attitude of the individual invites other people to work on or discuss a problem, and responsible individual's (responsible), the attitude will be responsible and ready to bear all the risks of everything he chooses (13.12). In-depth interviews with health workers show that respondents have with an incomplete immunization status that is mostly a sociocultural society or a belief that immunization is haram / should not be given to a baby, assuming immunization can lead to a child's fever, and not agreeing that the immunization is very important for is known child, it immunization is very important for The effects of fever children. perceived by children are temporary effect of the immunization because of the influence of immunization vaccines that are incorporated in the child's body, as well as the assumption of immunization, since vaccines given at immunization are not prohibited in religion. Respondents with social culture or beliefs tend to have attitudes negative towards immunization, so many children do complete not get basic immunization.

According the researcher's assumption the respondent's attitude in this research covers the comfort of the mother when the child is immunized, the mother's comfort after the child is immunized, the mother's attitude about the effect of immunization, religious view (halal / haram) of immunization. Factors that affect the number of respondents who have negative attitudes about immunization is a low knowledge of





immunization, the the lower knowledge of mother's immunization will contribute greatly to the formation of a bad attitude / about negative immunization, someone who already knows the truth about a thing then they will also have a positive attitude towards it, as well as immunization. The formation of this attitude is also inseparable from others who are considered important, mass media, emotional factors of the individual pengalamam about immunization. Preparation and implementation of complete basic immunization should be supported by a maximal role of health, cross-sector and community health officials, negative attitudes from the community about immunization need to be improved so that future generations can avoid certain infectious diseases, what can be done is to equate perceptions about the implementation of immunization with increasing public awareness of the importance of immunization, the side effects of immunization and the content of immunization vaccines given to infants. This is done in the hope that there is no presumption that immunization is not important, immunization is forbidden forbidden.

Healthservices

Based on the results of the study found more than half of respondents implementing complete basic immunization to get services that are less good.

The results of this study are in line with research conducted by Rahmawati stating that officers who are always present at the time of immunization services (95%) more

respondents who carry out complete basic immunization from officers who are not present during immunization schedule, as well as service officers who serving the mother and baby, the friendly officer (95%) is more likely to carry out complete basic immunization than the unhelpful officer while serving the respondent (mother and baby).

Health services are jointly or jointly organized within an organization to maintain and improve health, prevent and cure disease and restore the health of individuals, families, groups and even communities (14)

From the FGD results with the respondents in the Working Area of Puskesmas Pengambiran Padang City related to the health service that for some health workers who have given information and invite to complete the complete basic immunization, the respondent will come to posyandu, and for those who do not get the information and solicitation immediately sometimes forget to come to posyandu.

According to health officials who do not come during posyandu usually we visit one-one house after completion of posyandu whose baby is at home at the time of posyandu implementation.

According to the researcher's assumptions, the better the health service provided by the health worker then this will have an impact on the increase of health degree, as well as the immunization, the better the immunization service obtained by the parents and the baby, the greater the coverage of complete basic immunization, this is derived from the preparation of health



workers from the outset who approached the community by equating perceptions about complete basic immunization and not free from the high willingness of the community towards the implementation of complete basic immunization.

Multivariate Discussion

After double logistic regression test, the most related variable to the implementation of complete basic immunization is family support variable with p=0,001.

In multivariate analysis results can be concluded that if respondents get positive family support ithe implementation of complete basic immunization it will contribute to the achievement of the implementation of complete basic immunization of 8 times better.

The results of this study are relevant to Sri Enda Sitepu (2012) research, which states that there is family support (husbands, parents, in-laws and other relatives) to mothers in the form of getting information from families about basic immunization in children. Mom will feel that immunization is very important to boost her baby's This immunity. condition will certainly affect greatly the achievement of the expected immunization. Mothers who supported by family members to immunize their children tend to provide complete basic their immunization to children. otherwise mothers who are not supported by family members to immunize their children have a risk of not providing complete basic immunization to their children. The results of this study in accordance

with the theory that states that to realize the attitude into a real action required support factors or a condition that allows, among others, the facility. A positive mother's attitude to immunization should be confirmed by her husband and family, there is an easily accessible immunization facility, so that the mother may immunize her child (14)

From the FGD results with the respondents in the Working Areas of Puskesmas Pengambiran Padang obtained information that family support greatly affects the implementation of complete basic immunization because if the family does not approve then the mother also does not implement complete basic immunization. Mothers who give birth and have children receive decisions made by husband and family. According to health workers based on in-depth interviews, it is found that there is a myth in the family itself which states that immunization religion in immunization is not good for the baby, because it will cause the baby to become fever after immunization, implementation the immunization in infants is not done by the mother.

Officers provide a letter of agreement that can be accountable to the Department of Health if there is a family that does not conduct complete immunization.

support Family is the provided support by family members (husbands, wives, relatives) so that individuals who are given support feel that they are cared for, valued, getting help from meaningful people and have strong family ties with other family members (Lubis, Namora



Hasnida, 2009). Individuals who receive high family support will be more optimistic individuals in dealing with health and life issues and more skilled in meeting psychological needs (2)

According to the researchers' assumptions on the results of the relationship obtained in this study, in general mothers who do not get good family support caused by the myths dikeluarga itself that states that immunization is illegal in religion and immunization is not good for the baby, because it will cause the baby to be febrile after immunization, the implementation of immunization in infants is not performed by the mother.

CONCLUSION

The coverage of complete basic immunization has not reached 95% target because the system of implementation of complete basic immunization service which not yet optimal. It needs various efforts to achievement improve the complete immunization basic coverage, especially through optimizing the role and function ofhealth and cadre officers and optimizing partnerships with various parties.

REFERENCES

- Azwar, Your Human Attitudes Theory Measurement. Jakarta: Student Library 2008.
- Data and Information Center Ministry of Health RI. Immunization

- Situation in Indonesia. Jakarta 2016.
- Health MO. Indonesian Demographic And Health Survey (IDHS). Jakarta: Ministry Of Health: 2007.
- Lisnawati. Healthy Generation Through Immunization. Jakarta: Trans Info Media 2014; 5-14
- Law of the Republic of Indonesia. RI Law No. 36 Jakarta: Rineka Cipta 2009-5
- Ministry of Health RI. Indonesia Health Profile 2014. Jakarta: Ministry of Health RI; 2015
- Notoatmojo, Soekidjo. Health Promotion Theory & Applications. Publisher PT Adi Mahasatya: Jakarta 2010.
- Peshan Municipality of Padang City.
 Health Profile of Padang City
 2015. Padang: Padang City
 Health Office 2016.
- Priyoto. Changes in Health Behavior .. Yogyakarta: Graha Science 2015
- Rustikayanti, Suryamah. Perceptions About Booster Immunization in Mothers with Children Under 24 Months. 5 June 2017. STIKes Dharma Husada Bandung Jl. Jakarta Canal No. 75 Antapani Bandung 2017.
- Rizani. factor factors related to mother's behavior in giving immunization 2011. Diploma thesis, 2009 and also university
- Sugiyono. Quantitative Research Methods, Qualitative and R & D. Bandung: 2014; 21-28.
- Triana, vivi. Factors Associated with Complete Basic Immunization in Infants by 2015. University of Andalas. Padang.